

The Masses and Evolutionary State of the Stars in the Dwarf Nova SS Cygni¹

Martin A. Bitner and Edward L. Robinson

*Department of Astronomy, University of Texas at Austin, 1 University Station C1400,
Austin, TX, 78712*

`mbitner@astro.as.utexas.edu`

and

Bradford B. Behr

Code 7215, Naval Research Laboratory, Washington, DC 20375

ABSTRACT

The dwarf nova SS Cygni is a close binary star consisting of a K star transferring mass to a white dwarf by way of an accretion disk. We have obtained new spectroscopic observations of SS Cyg. Fits of synthetic spectra for Roche-lobe-filling stars to the absorption-line spectrum of the K star yield the amplitude of the K star's radial velocity curve and the mass ratio: $K_K = 162.5 \pm 1.0 \text{ km s}^{-1}$ and $q = M_K/M_{wd} = 0.685 \pm 0.015$. The fits also show that the accretion disk and white dwarf contribute a fraction $f = 0.535 \pm 0.075$ of the total flux at 5500 Å. Taking the weighted average of our results with previously published results obtained using similar techniques, we find $\langle K_K \rangle = 163.7 \pm 0.7 \text{ km s}^{-1}$ and $\langle q \rangle = 0.683 \pm 0.012$. The orbital light curve of SS Cyg shows an ellipsoidal variation diluted by light from the disk and white dwarf. From an analysis of the ellipsoidal variations we limit the orbital inclination to the range $45^\circ \leq i \leq 56^\circ$. The derived masses of the K star and white dwarf are $M_K = 0.55 \pm 0.13 M_\odot$ and $M_{wd} = 0.81 \pm 0.19 M_\odot$, where the uncertainties are dominated by systematic errors in the orbital inclination. The K star in SS Cyg is 10% to 50% larger than an unevolved star with the same mass and thus does not follow the mass-radius relation for Zero-Age Main-Sequence stars; nor does it follow the ZAMS mass/spectral-type relation. Its mass and spectral type are, however, consistent with models in which the core hydrogen has been significantly depleted.

Subject headings: binaries: close, stars: fundamental parameters, individual: SS Cygni

1. INTRODUCTION

As the brightest dwarf nova and one of the brightest cataclysmic variables of any kind, SS Cygni has been extensively observed and its properties are thought to be well established. Like all cataclysmic variables, it is a binary star consisting of a cool, relatively normal star and a white dwarf (Warner 1995). Indeed, it was one of the first cataclysmic variables shown to be a binary star – by Joy (1956), who found the orbital period to be $P_{orb} = 0.27624$ d, later revised to 0.27513 d by Stover et al. (1980) and Cowley et al. (1980). The spectral type of the cool star is K4-5 V (Smith & Dhillon 1998; Beuermann 2000). The K star fills its Roche lobe and is transferring mass through the inner Lagrangian point to an accretion disk around the white dwarf. During quiescence the disk has a low viscosity and accumulates mass, remaining cool and faint. When the surface density of the disk reaches a critical value, the viscosity increases and mass flows rapidly through the disk to the white dwarf, releasing gravitational energy. The disk becomes hot and luminous, causing the dwarf nova outburst (Lasota 2001). The outbursts recur roughly every 50 days (Cannizzo & Mattei 1992).

Standard techniques for measuring the masses of binary stars do not work for SS Cyg because only the spectrum of the K star is visible and the orbital light curve does not show eclipses. Previous measurements of the masses of the stars in SS Cyg have, therefore, been forced to rely on dubious assumptions about the properties of the accretion disk or about the structure of the K star (see section 5). In this paper we use a technique for measuring masses that was first developed for the black hole X-ray binaries and does not require these assumptions. The technique has three steps (eg, Casares 2006). First, the amplitude of the K star’s radial velocity curve K_K and the orbital period P can be combined to give the mass function

$$\frac{M_{wd} \sin^3 i}{(1+q)^2} = \frac{K_K^3 P}{2\pi G}, \quad (1)$$

where $q = M_K/M_{wd}$ is the ratio of the mass of the K star to the mass of the white dwarf and i is the orbital inclination. Second, the rotational velocity of the K star can be measured from the rotational broadening of its absorption lines. If the rotation period of the K star is tidally locked to the orbital period, the ratio of its projected rotational velocity $V_{rot} \sin i$ to K_K is a function of the mass ratio

$$\frac{V_{rot} \sin i}{K_K} = f(q). \quad (2)$$

¹Based on observations obtained with the Hobby-Eberly Telescope, which is a joint project of the University of Texas at Austin, the Pennsylvania State University, Stanford University, Ludwig-Maximilians-Universität München, and Georg-August-Universität Göttingen.

Third, the distorted K star causes ellipsoidal variations in the orbital light curve. The ellipsoidal variations give another relation between the mass ratio and the orbital inclination. Light from the accretion disk and white dwarf can distort the ellipsoidal variations, so it is also necessary to measure the amount of contaminating light from these other sources. The three relations together yield the dynamical masses of the stars and the dimensions of the system. The derived masses depend only weakly on assumed properties of the K star and the accretion disk.

We have obtained new spectroscopic observations of SS Cyg with the Hobby-Eberly Telescope (HET). We have measured K_K and q by fitting the spectrograms with model spectra calculated using our programs for synthesizing the rotationally broadened spectra of cool, Roche-lobe-filling stars (Bitner & Robinson 2006). The orbital light curve of SS Cyg has been measured by Kjurkchieva et al. (1999) and by Voloshina & Khruzina (2000) and shows a double-humped ellipsoidal variation diluted by light from the accretion disk. We analyze the ellipsoidal variations to place limits on the orbital inclination. Equipped with K_K , q , and i , we derive $M_K = 0.55 \pm 0.13 M_\odot$ and $M_{wd} = 0.81 \pm 0.19 M_\odot$. These masses are significantly smaller than the masses tabulated in the Ritter & Kolb (1998) catalog, $M_K = 0.704 M_\odot$ and $M_{wd} = 1.19 M_\odot$.

Section 2 of this paper describes the spectroscopic observations and section 3 describes how we measured K_K and q . In section 4 we describe our analysis of the ellipsoidal variations. Section 5 gives a detailed comparison of our results to earlier measurements and discusses the reasons for the differences. In the final section we derive the masses of the stars and discuss some implications of the lower masses.

2. OBSERVATIONS

The spectrograms of SS Cyg were acquired with the High Resolution Spectrograph (HRS) on the Hobby-Eberly Telescope (Tull 1998). We obtained 23 spectrograms in the 30 day interval from JD 2452084 to JD 2452114 (June and July 2001), each covering the wavelength range from 5300 Å to 7000 Å at a resolution $R = 30,000$. The exposure times were all 600 seconds, long enough to achieve the necessary signal-to-noise ratio without excessive spectral smearing from the changing radial velocity of the secondary star as it progressed around its orbit.

The heliocentric Julian dates at the midpoints of the exposures are given in Table 1. Figure 1 shows the eruption light curve of SS Cyg in 2001 from the American Association of Variable Star Observers (<http://www.aavso.org/data/lcg>). The horizontal bar plotted below

the light curve shows when we obtained our spectrograms. The figure also shows the eruption light curve in 1981 when it was observed by Hessman et al. (1984). Both sets of data were obtained when SS Cyg was in its quiescent state, but in 2001 the quiescent visual magnitude was $m_v = 12.0 - 12.1$, about 0.5 magnitudes fainter than in 1981.

The spectrograms were reduced using standard methods and standard IRAF packages. Two instrumental issues are of particular interest for the measurements of SS Cyg. The HRS has proved to have exceptionally good radial velocity stability, easily adequate for this project (Cochran et al. 2004). On the other hand, the HRS is a fiber-fed echelle spectrograph. Telescope tracking errors, variations in seeing, atmospheric dispersion, and other instrumental factors all limit the spectrophotometric accuracy of HRS spectrograms. Normalizing the continuum required care, especially because the free spectral range of the orders is rather small ($\sim 70 \text{ \AA}$).

As the optical/near-infrared spectrum of SS Cyg has been published many times (Echevarria et al. 1989; Friend et al. 1990; Martinez-Pais et al. 1994; North et al. 2001, 2002; Webb et al. 2002; Harrison et al. 2000), we show here only enough of the spectrum to demonstrate the rotational broadening of its absorption lines. Figure 2 shows the spectrum of SS Cyg near $6120 \text{ \AA}$. The spectrum shown is a time-averaged spectrum calculated by shifting the 23 individual spectrograms to SS Cyg’s systemic velocity and then adding them together. For comparison the figure also shows a synthetic spectrum for a single, non-rotating star with a K4-5 V spectral type calculated using the ATLAS9 stellar atmosphere program and the MOOG spectrum synthesis program. We show in the next section that the extra width of the absorption lines in SS Cyg is due to a combination of rotational broadening ($\sim 89 \text{ km s}^{-1}$), orbital smearing during the exposures ($< 26 \text{ km s}^{-1}$), and instrumental resolution of the spectrograph (8.2 km s^{-1}).

3. ANALYSIS OF THE SS CYGNI SPECTRUM

The Synthetic Spectra: We measured K_K , q , and the fraction of the flux contributed by the accretion disk and white dwarf by fitting the HET spectroscopy with synthetic spectra for cool, Roche-lobe-filling stars in close binary systems. We computed the synthetic spectra with our LinBrod program, which computes the spectra by summing wavelength-dependent, velocity-shifted, specific intensities over the distorted surface of the lobe-filling star (Bitner & Robinson 2006). The wavelength-dependent specific intensities are calculated using the ATLAS9 stellar atmosphere program and a modified version of the MOOG spectrum synthesis program (Snedden 1973; Kurucz 1993).

The synthetic spectra for the K star depend strongly on the orbital period, the orbital phase, the mass ratio, and the amplitude of its radial velocity curve; less strongly on the star’s metallicity and temperature; and only weakly on the orbital inclination, the gravity darkening exponent, and other input parameters to ATLAS9 and MOOG. As our primary goal was to determine K_K and q , we fixed the orbital period at its known value, we assumed the K star has a solar metallicity, and we adopted standard values for the gravity darkening exponent and the input parameters to ATLAS9 and MOOG. In preparation for the fits, we created large grids of synthetic spectra, two dimensions of the grid covering the possible ranges of K_K and q , and a third dimension covering 100 orbital phases. Separate grids were calculated for different values of the orbital inclination and effective temperature of the lobe-filling star. While we did not fit the temperature and inclination, the separate grids were used to determine the sensitivity of K_K and q to those parameters and to measure the dilution of the K star spectrum by flux from the accretion disk. To speed the calculation of Doppler shifts the synthetic spectra were placed on a logarithmic wavelength scale with a spacing of 1 km s^{-1} .

The synthetic spectra were convolved with the HRS instrumental profile and with an orbital smearing profile. We approximated the instrumental profile with a trapezoid. The full width at half maximum (FWHM) of the trapezoid was set equal to 8.2 km s^{-1} , the mean FWHM of a dozen isolated, narrow lines in spectrograms of the Th-Ar wavelength calibration lamp. We used a rectangle function to model orbital smearing. The width of the rectangle at each orbital phase was determined from $\Delta V = K_K(2\pi\Delta t/P)\cos 2\pi\phi$, where Δt is the length of the exposure and ϕ is the orbital phase. The zero point in phase is the time of superior conjunction of the white dwarf. For $K_K \approx 162 \text{ km s}^{-1}$ and a 10 minute exposure time the maximum orbital smearing is $\sim 26 \text{ km s}^{-1}$.

Determination of K_K and q : The synthetic spectra were fitted to the observed spectrograms by weighted least squares. As the spectrograms are greatly oversampled in wavelength, we were able to assign a weight to each wavelength from the internal scatter of the spectrograms. We fitted only those wavelength regions with several strong absorption features. There were seven of these regions between $5350 \text{ \AA}$ and $6470 \text{ \AA}$, each $25 \text{ \AA}$ to $57 \text{ \AA}$ wide and each mostly within the central 2/3 of a spectral order. As the regions are narrow, we normalized the continua of both the synthetic and the observed spectra to 1.0 by fitting straight lines to the continua, a different straight line for each region.

We fitted all the wavelength regions of all the spectrograms simultaneously, minimizing the reduced χ^2 of the fit to find the best values of K_K and q , and using the behavior of χ^2 in the neighborhood of the minimum to estimate the internal errors. There are three additional parameters in these fits: the flux contributed by the disk and white dwarf, the

systemic velocity γ , and the orbital phase offset $\Delta\phi$, where the $\Delta\phi$ is measured with respect to the Hessman et al. (1984) ephemeris: $T_0 = \text{HJD } 2,444,841.86899$ and $P = 0.27512973$ d. The flux from the disk and white dwarf was allowed to run free with a different, independent value for each of the seven wavelength regions. As the spectrograms cover the orbit of SS Cyg fairly uniformly, we decided to determine γ and $\Delta\phi$ separately from K_K and q . Initial estimates for γ and $\Delta\phi$ were made from fits based on initial estimates of K_K and q from Beuermann (2000) and Hessman et al. (1984). Holding these initial estimates for γ and $\Delta\phi$ fixed, we then re-fitted the spectra to obtain improved values for K_K , q , and the diluting flux from the accretion disk. With these improved values for K_K and q , we looped back and redetermined γ and $\Delta\phi$. The redetermined values of γ and $\Delta\phi$ did not differ significantly from the initial estimates, and so we were done.

The results are $K_K = 162.5 \pm 1.0 \text{ km s}^{-1}$, $q = 0.685 \pm 0.015$, $\Delta\phi = 0.0036 \pm 0.0010$, and $\gamma = -16.1 \text{ km s}^{-1}$. Figure 3 shows the contour plot of χ^2 as a function of K_K and q . The sign of $\Delta\phi$ is in the sense that the observed radial velocities occurred a few minutes later than predicted by the Hessman et al. (1984) ephemeris. We do not attach a standard deviation to our measurement of γ because we did not observe radial velocity standards, leaving the zero point of our radial velocity system uncertain. The γ velocity is, however, consistent with the heliocentric velocity found by North et al. (2002), $\gamma = -13.1 \pm 2.9 \text{ km s}^{-1}$.

To investigate the dependence of K_K and q on the effective temperature used to calculate the synthetic spectra of the K star, we derived K_K and q for two values of T_{eff} , 4560 K and 5000 K, the former corresponding to a K4 V spectral type and the latter to K2 V on the Tokunaga (2000) temperature scale. We note in passing that the local effective temperature varies considerably from point to point over the surface of the star. The temperatures we quote are the flux-weighted mean effective temperatures. The dependence on temperature is weak. The change of T_{eff} from 4560 K to 5000 K decreased K_K by only 0.9 km s^{-1} and increased q by 0.005. The values of K_K and q reported throughout this paper correspond to 4560 K.

The absorption-line profiles depend on the orbital inclination, so the derived values of K_K and q also depend on inclination. To investigate the strength of this dependence, we re-derived K_K and q for several values of i . The dependence is weak. Changing i from 40° to 50° changed K_K by less than 0.5 km s^{-1} and q by about ~ 0.001 . Since the absorption-line profiles depend on the orbital inclination, they can be used to determine the inclination, at least in principle (Shahbaz 1998). In practice the dependence is so weak that fits to the spectra do not give reliable inclinations (Bitner & Robinson 2006). Indeed, for SS Cyg our fits would suggest an inclination near or lower than 30° , an impossibly low value since it would imply a white dwarf mass much greater than the Chandrasekhar limit. As a result,

we do not determine the inclination this way.

Determination of the Flux from the Accretion Disk and White Dwarf: The effect of extra flux from the accretion disk and white dwarf is to dilute the absorption lines in the spectrum of the K star. Since the depth of the K star’s absorption lines depends on its spectral type, the measured amount of flux from the disk and white dwarf depends on the spectral type and temperature adopted for the star. The dependence is strong. For $T_{eff} = 4560$ K the fits yielded $f = 0.49$, where f is the fraction of the flux at $5500 \text{ \AA}$ coming from the disk and white dwarf; and for $T_{eff} = 5000$ K the fits yielded $f = 0.260$. To measure f we must, therefore, have a more precise value for the temperature of the K star than we have needed up to now. We obtain this temperature in two steps: First we determine the precise spectral type of the K star, then we assign a temperature to that spectral type.

With the exception of Joy (1956), who classified the late type star as dG5, and of Martinez-Pais et al. (1994), who classified it as K2-K3, there is a broad consensus that the late-type star has a K4 V or K5 V spectral type (Stover et al. 1980; Cowley et al. 1980; Walker 1981; Bailey 1981; Wade 1982; Friend et al. 1990; Harrison et al. 2000; Webb et al. 2002; North et al. 2002). The spectral type does not depend on the method nor the wavelength region used for the measurement. Neither spectral type is preferred over the other. Thus, in their reviews of cataclysmic variable secondary stars Smith & Dhillon (1998) assigned a spectral type of K5 V to the secondary of SS Cyg while Beuermann (2000) assigned a spectral type of K4 V. We adopt K4.5 V with a half subclass uncertainty. The surface of the K star is likely to have regions with much lower temperatures (starspots) (Webb et al. 2002). Their effect on the spectral type is assumed to be negligible.

From Table 7.6 in Tokunaga (2000) the temperature corresponding to spectral type K4.5 V is 4450 K, while from Table 15.7 in Drilling & Landolt (2000), the temperature is 4480 K. Allowing for a half spectral-subclass uncertainty and a possible systematic error of 100 K due to starspots, heating by irradiation, or an incorrect temperature scale, we take $T_{eff} = 4470 \pm 140$ K as the best estimate of the temperature of the K star.

As noted above, for $T_{eff} = 4560$ K we find $f = 0.49$, and for $T_{eff} = 5000$ K we find $f = 0.260$. Extrapolating somewhat, we find $f = 0.535 \pm 0.075$ for $T_{eff} = 4470 \pm 140$ K. We note that errors in assumed metallicity can affect the line depths in our model LinBrod spectra and thus the derived flux contribution from the accretion disk and white dwarf. To quantify the magnitude of the effect, we compared results for LinBrod model spectra of solar metallicity and $[M/H] = -0.5$. This is a much larger metallicity error than we expect is reasonable. Even so, the derived flux contribution from the white dwarf and accretion disk only drops by 20-25% for the lower metallicity spectra. We measured the disk flux in each of the seven wavelength regions separately. While the results are rather noisy, the flux

contributed by the disk is roughly 20% less at 6400 Å than at 5500 Å. Other spectroscopic measurements of the disk fraction are listed in Table 2 for comparison. With the possible exception of the measurement by North et al. (2002) there is surprisingly good agreement among them.

An External Check on the Results: As a final external check on our results one of us (BBB) re-reduced the spectrograms of SS Cyg with the Figaro package and re-analyzed them with the LINFOR spectral synthesis package (Baschek et al. 1966; Lemke 1997, private communication), both of which are genetically unrelated to the IRAF and LinBrod packages. Using LINFOR we generated a model spectrum for the secondary, including a spherically-symmetric rotation profile with $v \sin i = 100$ km/s, and cross-correlated this model with each observed spectrum to get initial radial velocities. We then adjusted the model rotational velocity and the abundances of the major line-forming atomic species to minimize the residuals between the model and the observations. Finally, we used the model to identify regions of continuum and pseudo-continuum, and adjusted the adopted continuum levels for the observed spectra accordingly. The entire sequence was repeated until the results converged. We list the resulting radial velocities in Table 1. The rotational velocity varied from about 84 to 94 km s⁻¹, depending on orbital phase, with a mean near $V_{rot} \sin i = 89$ km s⁻¹. This method does not yield a reliable mass ratio because of the spherical approximation, and it does not yield a measure of the dilution because the dilution is mimicked by a reduction of metallicity; but the radial velocities should be unbiased (Bitner & Robinson 2006). A fit to the velocities yielded $K_K = 162.8$ km s⁻¹ in excellent agreement with the LinBrod results.

4. LIGHT CURVE ANALYSIS AND THE ORBITAL INCLINATION

The Source of the Orbital Variations: Kjurkchieva et al. (1999) and Voloshina & Khruzina (2000) have independently published multicolor mean orbital light curves for SS Cyg and the two sets of light curves agree to ~ 0.01 mag. We use the Voloshina & Khruzina (2000) data for our analysis because they give the variances of the individual points in the light curves. The mean V-band light curve from Voloshina & Khruzina (2000), reproduced in Figure 4, shows a distinctive double-humped modulation with minima near orbital phases $\phi = 0.0$ and 0.5 , and unequal maxima near $\phi = 0.25$ and 0.75 , where phase zero is again defined to be superior conjunction of the white dwarf. We attribute the double humped modulation primarily to ellipsoidal variations of the K star. The best justification for this interpretation is the quantitative agreement we find between the observed light curves and synthetic light curves based on ellipsoidal variations. The double humps have occasionally been attributed to spots on the white dwarf, caused perhaps by accretion onto magnetic

poles (Kjurkchieva et al. 1999). Since the humps remain at the same orbital phases, this interpretation would require the that rotation period of the white dwarf be identical to the orbital period. There is no evidence that SS Cyg has a magnetic field strong enough to lock on to the K star and enforce synchronism (ie, it is not an AM Her star) and, on the contrary, there is evidence that the rotation period of the white dwarf is just a few seconds (Mauche 2004). Spots on the white dwarf cannot then account for the double humped modulation.

A pure ellipsoidal variation produces two symmetric humps in a light curve, but in SS Cyg the hump at $\phi = 0.75$ has a larger amplitude than the hump at $\phi = 0.25$. The multicolor light curves published Kjurkchieva et al. (1999) and Voloshina & Khruzina (2000) show that difference between the two humps becomes progressively greater at shorter wavelengths, demonstrating that the extra flux in the larger hump is bluer than the flux in the pure ellipsoidal variations. The disks in cataclysmic variables generally have a hot spot on their edge where they are hit by a stream of gas transferred from the lobe-filling star, and the hot spot often produces a single hump in the orbital light curve peaking at phases between 0.7 and 0.9 (Warner 1995). Since the extra blue flux in SS Cyg is greatest near phase 0.75, we attribute the extra blue flux to a hot spot on the disk. Our interpretation of the light curve is, thus, similar to that of Voloshina & Khruzina (2000).

A Model for the Light Curve: Our goal is to measure the orbital inclination of SS Cyg from the amplitude of the ellipsoidal variations. The ellipsoidal variations are, however, diluted by unmodulated light from the accretion disk. This complicates the task because the ellipsoidal variations of binaries with low inclinations and low dilution are nearly identical to those of binaries with higher inclinations and greater dilution. Voloshina & Khruzina (2000) attempt to break the degeneracy between inclination and dilution by fitting multicolor light curves. For this method to work, the spectral energy distributions of all sources of flux must be known. They assumed that all the sources emit black body radiation. This is a poor assumption for the optical flux from the quiescent accretion disks of dwarf novae (Vrielmann & Baptista 2002; Saito & Baptista 2006) and is not really valid even for the flux from the two stars, making the orbital inclination determined from the multicolor fits unreliable.

The degeneracy between the orbital inclination and the disk flux can also be broken if the disk flux can be independently measured. The disk flux then becomes an input parameter, not a fitted parameter, and we can dispense with theoretical models for the disk flux and spectral energy distribution. From the fits to the spectrum described in section 3, we do know the V-band flux from the disk, so this is the method we will use to break the degeneracy.

The computer program we use to calculate light curves for SS Cyg is an updated and improved version of the program we have used to calculate the light curves of black hole

X-ray binaries (Ioannou et al. 2004). Although the program accepts many parameters and can synthesize light curves for complicated accretion geometries, just three components are needed to fit the light curve of SS Cyg:

- **The Ellipsoidal Variations:** We use Roche geometry to calculate the shape and local effective gravity of the lobe-filling K star. The local effective temperature $T_{eff}(local)$ is calculated from the gravity darkening law $T_{eff}(local) \propto T_{eff}|g|^\beta$, where g is the local gravity, and β is the gravity-darkening exponent. According to Claret (2000a) $\beta \approx 0.10$ for main-sequence stars with temperatures near 4500 K. The amplitude of the ellipsoidal variations depends only weakly on β . The angle-averaged fluxes in specific filters are calculated by integrating Kurucz model spectra over the filter response functions. We adopt the Claret (2000b) limb darkening law to convert the fluxes to the local specific intensities, which are then integrated over the surface of the star to produce the observed brightness.
- **Constant Flux from the Disk:** The main body of the accretion disk is uneclipsed and just adds constant extra flux to the light curve. Any contribution from the white dwarf is subsumed in the disk contribution.
- **Variable Flux from a Bright Spot on the Disk:** The disk has a bright spot painted on its outer edge. The spot is visible only when on the side of the accretion disk facing the observer and its brightness is modulated by geometric foreshortening. The properties of the spot are specified by its position and extent along the disk edge and by its brightness.

The Fits to the Light Curve: Although Voloshina & Khruzina (2000) published light curves in the U, B, and V bands, we fit only the V-band light curve because the distorting effect of the bright spot is least in that filter and because the V band is the only one for which our spectroscopy gives a measurement of the diluting flux from the disk. For all fits we use $K_K = 162 \text{ km s}^{-1}$ and $q = 0.69$. The deduced orbital inclinations are essentially independent of K_K , and changing the mass ratio by $\Delta q = 0.01$ changes the deduced orbital inclinations by just 0.5° . The errors introduced by uncertainties in K_K and q are, therefore, small compared to other sources of error. Since we are fitting just the V-band light curve, the deduced inclinations are almost independent of the adopted effective temperature, with just a second-order dependence through the limb-darkening and the gravity darkening coefficient. For convenience internal to the computer program we used $T_{eff} = 4600 \text{ K}$, not 4470 K. Numerical tests showed that the ellipsoidal light curves and the derived inclinations are not significantly changed by the difference in temperature.

We proceeded by choosing a specific value for the fraction of the V-band flux coming from the disk (at $\phi = 0.25$) and then fitting the synthetic light curve to the data by adjusting the remaining parameters to minimize the χ^2 of the fit. The four physically-significant parameters are the orbital inclination, the location of the spot on the edge of the disk, the extent of the spot along the edge, and the total V-band flux from the spot. There are also two nuisance parameters: the offset in orbital phase between the synthetic and observed light curves, and an overall scale factor. We repeated the fit for many different values of the fraction of the flux coming from the disk, producing a table of orbital inclinations and standard deviations as a function of the disk flux.

Figure 4 shows a synthetic light curve over-plotted on the observed light curve. The synthetic light curve corresponds to an orbital inclination of 52° with 56% of the light coming from the accretion disk and a phase offset of $\Delta\phi = 0.01$. The amplitude of the ellipsoidal variations – and therefore the deduced orbital inclination – is determined almost entirely by the amplitude of the hump at phase 0.25, whereas the properties of the spot are determined almost entirely by the hump at phase 0.75. As a result, the orbital inclinations depend only weakly on the spot model. Figure 5 shows the deduced orbital inclination as a function of the fraction of the V band flux coming from the accretion disk. The best fit inclinations are shown by the solid line extending from the lower-left to the upper right in the diagram. The dashed lines above and below the solid line show the one standard deviation confidence limits for the fits.

The Orbital Inclination: In the previous section we found that the fraction of the V band flux coming from the disk was $f = 0.535 \pm 0.075$. The vertical solid line in Figure 5 corresponds to $f = 0.535$ and the two vertical dashed lines to $f = 0.46$ and $f = 0.61$. The allowed values of the disk flux and orbital inclination lie in the region enclosed by the four dashed lines in the figure. Rounding the artificially sharp corners of the region somewhat, we find that the orbital inclination of SS Cyg lies in the interval $45^\circ \leq i \leq 56^\circ$. The range of possible inclinations is rather wide. The width is dominated by the uncertainty in f , which traces back to the uncertainty in the temperature of the K star determined from its spectral type; other sources of error are minor contributors.

5. COMPARISON TO PREVIOUS RESULTS

As the literature on SS Cyg is replete with discrepant measurements of K_K , q , i and the masses of its components, and as our measurements lead to yet another set of masses, it is appropriate to give a critical review of the literature, comparing our results to the previous measurements and analyzing the reasons for the differences.

Radial Velocity Curve of the K Star: Table 3 lists all the published measurements of K_K . The amplitudes fall into three groups. The first consists of early measurements yielding amplitudes near 120 km s^{-1} . In this group the spectrograms were recorded on photographic plates and velocities were measured using traditional techniques. Beginning with Stover et al. (1980) the spectrograms were obtained with digital detectors and the velocities were usually measured by cross-correlating or directly fitting a template spectrum to the spectrograms. Initially controversial, Stover’s measurement was the first of a group made between 1980 and 1990 yielding amplitudes near 155 km s^{-1} . There are two main reasons for the difference between these and the earlier results. First, the absorption lines from the K star are broadened by rotation and veiled by continuum emission from the accretion disk, making line positions difficult to measure on photographic plates. Second, heating of the K star by irradiation from the accretion disk during dwarf nova outbursts distorts the radial velocity curve, changing the measured amplitude (Robinson et al. 1986). While the distortion has not been fully modeled theoretically, the presence of the distortion is amply demonstrated by observations (Martin et al. 1989; Echevarria et al. 1989). Observers have since become careful to measure the radial velocity curve of the K star during quiescence, although Robinson et al. (1986) suggested that there might be enough irradiation to bias the measured value of K_K significantly even during quiescence.

The radial velocity amplitudes measured by Martinez-Pais et al. (1994) and North et al. (2002), 162.5 ± 3 and $165 \pm 1 \text{ km s}^{-1}$ respectively, and now $162.5 \pm 1.0 \text{ km s}^{-1}$ by us, comprise the third group.² While the difference between these amplitudes and the $\sim 155 \text{ km s}^{-1}$ amplitudes of the second group is small, it is statistically significant. We attribute the difference to a real change in the SS Cyg system. From Cannizzo & Mattei (1992) the mean quiescent visual magnitude of SS Cyg between 1896 and 1992 was $m_v = 11.8 - 11.9$. The quiescent magnitude fluctuated by a few tenths of a magnitude about the mean and appeared to show a ~ 0.1 magnitude secular decrease over the 100 year interval. Hessman et al. (1984) observed SS Cyg when its quiescent magnitude was $m_v = 11.5 - 11.6$, about 0.3 magnitude brighter than the historical average (see Figure 1). AAVSO light curves (<http://www.aavso.org/data/lcg>) show that SS Cyg was at $m_v = 11.9 - 12.0$ when observed by Martinez-Pais et al. (1994), at $m_v \approx 12.0$ when observed by North et al. (2002), and from Figure 1 at $m_v = 12.0 - 12.1$ when observed by us. Succinctly, the recent observations were obtained when SS Cyg was fainter than its historical average and much fainter than when observed by Hessman et al. (1984). Following Cannizzo & Mattei (1992) and Warner (1995), we attribute a lower quiescent brightness to a lower accretion luminosity, which implies lower

²Martinez-Pais et al. (1994) measured $162.5 \pm 3 \text{ km s}^{-1}$ from their own data but then adopted $158 \pm 3 \text{ km s}^{-1}$ from Robinson et al. (1986) in the rest of their paper.

irradiation of the K star and less distortion of its radial velocity curve. As a result, the values of K_K in the third group should be a better representation of the true motion of the K star.

The Mass Ratio: As the white dwarf in SS Cyg is a minor contributor to the flux at optical and infrared wavelengths, its radial velocity curve cannot be directly measured and the mass ratio must be deduced by other means. Lacking a better alternative it has sometimes been assumed that the wings of the emission lines, which come from high-velocity gas near the white dwarf, can serve as a proxy for the white dwarf and track its orbital motion. Measurements of the amplitude K_{em} of the emission-line radial velocity curve of SS Cyg are shown in Table 3. The weighted average of the amplitudes measured with electronic detectors is $K_{em} \approx 92 \text{ km s}^{-1}$. Taking $K_K \approx K_{abs} \approx 152 - 165 \text{ km s}^{-1}$, one finds $q = M_K/M_{wd} = 0.56 - 0.60$, in accord with the value tabulated by Ritter & Kolb (1998). Although one of us has previously argued that emission lines can yield good qualitative estimates of the mass ratios (Robinson 1992), more recent work has demonstrated that emission-line velocities can yield biased or simply wrong mass ratios and should not be used for accurate work [eg, Welsh et al. (1993)].

We have measured the mass ratio from the rotational broadening of the absorption lines in the spectrum of the K star. This technique has been widely used to measure the masses of black holes in black hole binary stars and, if applied with care, gives reliable mass ratios (Casares 2006). Cowley et al. (1980) used line broadening as a consistency check on other ways to measure the mass ratio of SS Cyg but Martinez-Pais et al. (1994) were the first to use line broadening as the fundamental way to measure the mass ratio, finding $q = 0.62 \pm 0.05$. Their measurement was, however, biased by the adoption of a Gray (1992) rotational broadening profile, which is only appropriate for spherical stars, and by the use of approximate formulae for converting from rotational velocity to mass ratio. North et al. (2002) also measured the mass ratio from the rotational broadening of the K star’s spectrum. Their method, outlined in North et al. (2000), is similar to ours and like ours should be free of bias. Their result, $q = 0.68 \pm 0.02$, is nearly identical to ours.

The Orbital Inclination: SS Cyg does not eclipse, so the orbital inclination must be measured indirectly. One way has been to assume the K star has some known property and then work backwards to the orbital inclination. If one assumes that the K star follows the mass/spectral type relation for main-sequence stars, its mass is $0.7 - 0.8 M_\odot$ and the inferred orbital inclination is $38^\circ - 40^\circ$ (Cowley et al. 1980; Walker 1981). It is more common to assume that the K star follows the main-sequence mass/radius relation. There are many variants of this relation but all lead to orbital inclinations in the range $35^\circ - 45^\circ$ (Kraft 1962; Kiplinger 1979; Stover et al. 1980; Hessman et al. 1984; Hessman 1988; Friend et al. 1990). The properties of lobe-filling stars can, however, depart greatly from main-sequence

relations because of nuclear evolution before the onset of mass transfer and the loss of thermal equilibrium while mass is being transferred. The theoretical calculations by Kolb et al. (2001) show that lobe-filling stars with K4 – K5 spectral types can have masses anywhere in the range $0.42 - 0.80 M_{\odot}$. Baraffe & Kolb (2000) show that the radii of stars that have used up some of the hydrogen in their cores and are transferring mass can be larger than the radii of normal main-sequence stars with the same mass by 50% or more. The observational data collected by Smith & Dhillon (1998) and Beuermann (2000) support these calculations.

There have been several attempts to estimate the orbital inclination from properties of the accretion disk. Comparing the profile of the $H\beta$ emission line in SS Cyg to theoretical profiles for a thin disk, Cowley et al. (1980) derived an inclination between $30 - 40^{\circ}$. If the inner radius of the accretion disk is only slightly greater than the radius of the white dwarf and if the white dwarf obeys the usual white dwarf mass/radius relation, one can use the inner radius to deduce the mass of the white dwarf and again work backwards to the orbital inclination. Cowley et al. (1980) estimated the radius from the maximum velocity extent of the wings of the Balmer emission lines, finding $i \approx 46^{\circ}$. Giovannelli et al. (1983) assumed that the periods of dwarf nova oscillations (DNOs) are equal to the orbital periods of gas at the inner edge of the accretion disk. At that time the DNOs of SS Cyg had been observed to range from 7.3 s to 10.9 s, from which Giovannelli et al. (1983) derived $i = 40^{\circ} \pm 2^{\circ}$. Doppler tomograms demonstrate, however, that the velocity fields of accretion disks, including the accretion disk of SS Cyg, are complicated (Robinson et al. 1993; North et al. 2001), and the velocities are likely to be especially complicated near the white dwarf where magnetic fields may play a role; and the periods of DNOs are only indirectly related to Keplerian periods (Warner 2004; Kluźniak et al. 2005), all of which precludes giving much weight to inclinations derived from properties of the accretion disk.

The analysis of orbital light curves generally yield reliable inclinations. This is clearly true for eclipsing binaries but extensive observations of the black hole binaries have shown that this is also true for orbital inclinations derived from the ellipsoidal variations of Roche-lobe-filling stars (Charles & Coe 2006). Ellipsoidal variations are produced primarily by the geometric distortions of the star, which, since the star fills its Roche lobe, are a function only of the mass ratio. Large amplitude ellipsoidal variations imply a large inclination, low amplitude a low inclination. The rapid flickering in the light curve of SS Cyg and its awkwardly-long orbital period vitiated earlier attempts to measure its ellipsoidal variations (eg, Honey et al. 1989). The variations are, however, present and have been observed at both ultraviolet and optical wavelengths (Bartolini et al. 1985; Voloshina 1986; Lombardi et al. 1987; Bruch 1990; Kjurkchieva et al. 1999; Voloshina & Khruzina 2000). The ellipsoidal variations of SS Cyg are diluted by flux from the accretion disk and this dilution must be included in the models if fits to the light curves are to give reliable inclinations. Voloshina & Khruzina

(2000) dealt with this by assuming all components of the SS Cyg system radiate like black bodies, finding $i = 51^\circ - 56^\circ$, but their assumption is questionable, raising concerns about the reliability of their result. We have estimated the disk contribution externally, from the dilution of the K star’s spectrum. Consequently we were able to use a far simpler model for the light curve, which leads in turn to a more robust estimate of the orbital inclination. Nevertheless, the range of inclinations we find, $45^\circ \leq i \leq 56^\circ$, is not incompatible with the range of inclinations found by Voloshina & Khruzina (2000). The qualitative reason why orbital inclinations derived from the ellipsoidal variations is high is that the observed ellipsoidal variations have a large amplitude even though they are heavily diluted by flux from the accretion disk. Unless the measured orbital light curves incorrectly represent the true light curve, it is difficult to escape the higher inclination.

6. DISCUSSION

The Masses of the Stars: From fits of synthetic spectra to spectrograms of SS Cyg obtained with the HET we have found the amplitude of the K star’s radial velocity curve to be $K_K = 162.5 \pm 1.0 \text{ km s}^{-1}$ and the ratio of the masses of the two stars to be $q = M_K/M_{wd} = 0.685 \pm 0.015$. From an analysis of the ellipsoidal variations in the orbital light curve plus an estimate of the dilution of the ellipsoidal variations by flux from the accretion disk, we have limited the orbital inclination to the range $45^\circ \leq i \leq 56^\circ$.

Following our discussion in section 5, we see no reason not to accept the measurements of K_K by Martinez-Pais et al. (1994) and North et al. (2002) and the measurement of q by North et al. (2002). The best estimates of K_K and q are, then, the weighted average of their measurements and ours, $\langle K_K \rangle = 163.7 \pm 0.7 \text{ km s}^{-1}$ and $\langle q \rangle = 0.683 \pm 0.012$. Adopting these values and taking $45^\circ \leq i \leq 56^\circ$, we find masses $M_K = 0.55 \pm 0.13 M_\odot$ and $M_{wd} = 0.81 \pm 0.19 M_\odot$. The uncertainties in the masses are dominated by the uncertainty in the orbital inclination, which is itself dominated by systematic errors in the temperature of the K star. As a result, it is best to interpret these values as the ranges of possible masses and the midpoints of the ranges, not the best masses and their standard deviations. A Monte-Carlo error analysis suggests that lower masses within the range are somewhat more likely than higher masses. The masses of the K star and the white dwarf cannot be chosen independently from within their ranges as they are strongly constrained by the mass ratio $M_K/M_{wd} = 0.683 \pm 0.012$. Kepler’s third law now yields the separation of the centers of mass of the two stars $a = (1.36 \pm 0.11) \times 10^{11} \text{ cm}$, where higher values of a correspond to higher masses.

The Evolutionary State of the K Star: If the mean radius of a Roche lobe is R_L , then

R_L/a is a function only of the mass ratio, $R_L/a = F(q)$. Since SS Cyg is a mass transfer system, the radius of the K star R_K equals the radius of its Roche lobe and can be calculated from $R_K/a = F(q)$. Adopting the Eggleton (1983) relation for $F(q)$ and using our measurement of a and M_K as a function of i , we find $R_K/M_K = 1.07$ at ($i = 45^\circ$, $M_K = 0.68$), increasing to $R_K/M_K = 1.47$ at ($i = 56^\circ$, $M_K = 0.42$), where the units of R and M are solar radius and solar mass.

From the theoretical models of Chabrier & Baraffe (1997) the mass-radius relation for zero age main sequence (ZAMS) stars with masses between $0.4 M_\odot$ and $0.6 M_\odot$ is $R = 0.95M$, or $R/M = 0.95$, where here also the units are solar radius and solar mass. Thus, the K star in SS Cyg is 10% to 50% larger than a single unevolved star with the same mass – it does not obey the mass-radius relation for unevolved main-sequence stars.³ Nor does the K star follow the ZAMS mass/spectral-type relation for Roche-lobe-filling stars. Kolb et al. (2001) have calculated theoretical evolutionary models for the secondary stars of mass-transfer binaries. They find that an unevolved lobe-filling star with spectral type K4-5 (their “standard sequence”) should have a mass $\sim 0.2 M_\odot$ greater than the mass of the K star in SS Cyg. Finally, according to Howell et al. (2001) an unevolved lobe-filling star in a binary with SS Cyg’s 6.6 hr orbital period should have a mass $\sim 0.2 M_\odot$ greater than that of the K star in SS Cyg.

The Kolb et al. (2001) models do, however, predict lower masses for K4-5 stars in which core hydrogen is depleted but not exhausted. If the mixing length is maintained at $\alpha = 1.0$, the depletion must be extreme to match SS Cyg, $X_c = 4 \times 10^{-4}$; but if the mixing length is increased to $\alpha = 1.9$, the depletion can be more moderate. We interpret our results, therefore, as showing that the K star in SS Cyg has significantly depleted its hydrogen.

Harrison et al. (2004) showed that the CO absorption bands in the infrared spectrum of SS Cyg are anomalously weak and Harrison et al. (2005) found weak CO bands in 9 of 12 other cataclysmic variables they observed. They attribute the weak CO bands to a low C abundance and note that C can be depleted by CNO processing. While the CNO cycle can indeed deplete C, all that is really needed is to process C to N; the full CNO cycle is not required. In either case C should not be depleted in normal main-sequence stars, so the weak CO bands are evidence for nuclear evolution. We take this as supporting evidence that the K star in SS Cyg is not an unevolved main-sequence star.

³If we were to discard the orbital inclination determined from the ellipsoidal variations and insist that the K star obeys the Chabrier & Baraffe (1997) ZAMS mass-radius relation, we would find $M_K = 0.81 M_\odot$, $M_{wd} = 1.18 M_\odot$, and $i = 42^\circ$. These are stringent upper limits on the masses and lower limit on the orbital inclination.

The Visual Magnitude of SS Cyg: To calculate the expected V magnitude of SS Cyg, we begin with the absolute visual magnitude of a normal K4.5 V star, which from Table 15.7 of Drilling & Landolt (2000) is $M_V = 7.19$. The radius of the K star in SS Cyg is 10% to 50% larger than normal. Scaled by its surface area the K star’s absolute magnitude should be greater than normal by 0.21 to 0.88 magnitude, or $M_{V_K} = 6.64 \pm 0.34$. The trigonometric parallax of SS Cyg is 6.02 ± 0.46 milliarcseconds, leading to a distance modulus $\Delta m = 6.10 \pm 0.16$. The apparent visual magnitude of the K star alone should then be $V_K = 12.74 \pm 0.37$. This must be augmented by the flux from the accretion disk and white dwarf, which we have measured to be a fraction $f = 0.535 \pm 0.075$ of the total V-band flux. The extra flux raises the expected V magnitude of the entire system to $V = 11.92 \pm 0.40$, which is consistent with the visual magnitude $m_v = 12.0 - 12.1$ measured by the AAVSO at the time of our observations.

We thank the American Association of Variable Star Observers for the data used to make Figure 1, Carlos Allende Prieto for help with HET HRS data reduction, Robert A. Wittenmyer for performing an independent fit of a spectroscopic orbit to the velocities listed in Table 1, William F. Welsh for comments on a draft of this paper and for performing the Monte-Carlo error analysis, and Chris Sneden for help using and modifying the MOOG program. This research was supported by NSF grant AST-0206029. MAB acknowledges support from NSF grant AST-0607312. This research has made use of NASA’s Astrophysics Data System (ADS) Bibliographic Services. The Hobby-Eberly Telescope (HET) is a joint project of the University of Texas at Austin, the Pennsylvania State University, Stanford University, Ludwig-Maximilians-Universität München, and Georg-August-Universität Göttingen. The HET is named in honor of its principal benefactors, William P. Hobby and Robert E. Eberly.

REFERENCES

- Bailey, J. 1981, MNRAS, 197, 31
- Baraffe, I., & Kolb, U. 2000, MNRAS, 318, 354
- Bartolini, C., Guarnieri, C., Lolli, M., Piccioni, A., Giovannelli, F., Gaudenzi, S., & Lombardi, R. 1985, in Multifrequency Behaviour of Galactic Accreting Sources, Ed. by F. Giovannelli (Frascati: CNR, Istituto di Astrofisica), p.50
- Baschek, B., Holweger, H., & Traving, G. 1966, AAHam, 8, 26B
- Beuermann, K. 2000, New Astron. Rev., 44, 93

- Bitner, M. A., & Robinson, E. L. 2006, *AJ*, 131, 1712
- Bruch, A., 1990, *Acta Aston.*, 40, 369
- Cannizzo, J, K., & Mattei, J. A. 1992, *ApJ*, 401, 642
- Casares, J. 2006, in “The Many Scales of the Universe - JENAM 2004 Astrophysics Reviews” eds. J. C. del Toro Iniesta et al. (Kluwer Academic Publishers)
- Chabrier, G., & Baraffe, I. 1997, *A&A*, 327,1039
- Charles, P. A., & Coe, M. J. 2006, in “Compact Stellar X-Ray Sources” eds. W. Lewin & M. van der Klis (Cambridge: Cambridge University Press) p. 215
- Cochran, W. D., et al. 2004, *ApJ*, 611, L133
- Claret, A. 2000, *A&A*, 359, 289
- Claret, A. 2000, *A&A*, 363, 1081
- Cowley, A. P., Crampton, D. & Hutchings, J. B. 1980, *ApJ*, 241, 269
- Drilling, J. S., & Landolt, A. U. 2000, in *Allen’s Astrophysical Quantities*, ed. A.N. Cox (4th ed.;New York: Springer), 381
- Echevarria, J., Diego, F., Tapia, M., Costero, R., Ruiz, E., Salas, L., Gutierrez, L., & Enriquez, R. 1989, *MNRAS*, 240, 975
- Eggleton, P. P. 1983, *ApJ*, 268, 368
- Friend, M. T., Martin, J. S., Smith, R. C., & Jones, D. H. P. 1990, *MNRAS*, 246, 654
- Giovannelli, F., Gaudenzi, S., Rossi, C., & Piccioni, A. 1983, *Acta Astron.*, 33, 319
- Gray, D. F. 1992, *The Observation and Analysis of Stellar Photospheres* (Cambridge: Cambridge Univ. Press)
- Harrison, T. E., McNamara, B. J., Szkody, P., McArthur, B. E., Benedict, G. F., Klemola, A. R., & Gilliland, R. L. 1999, *ApJ*, 515, L93
- Harrison, T. E., McNamara, B. J., Szkody, P., & Gilliland, R. L. 2000, *ApJ*, 120, 2649
- Harrison, T. E., Osborne, H. L., & Howell, S. B. 2004, *AJ*, 127, 3493
- Harrison, T. E., Osborne, H. L., & Howell, S. B. 2005, *AJ*, 129, 2400

- Hessman, F. V., Robinson, E. L., Nather, R. E., & Zhang, E. H. 1984, *ApJ*, 286, 747
- Hessman, F. V. 1988, *A&AS*, 72, 515
- Honey, W. B., Bath, G. T., Charles, P. A., Whitehurst, R., Jones, D. H. P., Echevarria, J., Arvalo, M. J., Solheim, J.-E., Tovmassian, G., & Takagishi, K. 1989, *MNRAS*, 236, 727
- Howell, S. B., Nelson, L. A., & Rappaport, S. 2001, *ApJ*, 550, 897
- Ioannou, Z., Robinson, E. L., Welsh, W. F., & Haswell, C. A. 2004, *AJ*, 127, 481
- Joy, A. H. 1956, *ApJ*, 124, 317
- Kiplinger, A. L. 1979, *AJ*, 84, 655
- Kjurkchieva, D., Marchev, D., & Ogloza, W. 1999, *Ap&SS*, 262, 53
- Kluźniak, W., Lasota, J.-P., Abramowicz, M. A., & Warner, B. 2005, *A&A*, 440, 25
- Kolb, U., King, A. R., & Baraffe, I. 2001, *MNRAS*, 321, 544
- Kraft, R. P. 1962, *ApJ*, 135, 408
- Kurucz, R. 1993, *ATLAS9 Stellar Atmosphere Programs and 2 km/s grid*. Kurucz CD-ROM No. 13 (Cambridge: Mass.: Smithsonian Astrophysical Observatory)
- Lasota, J.-P. 2001, *New Astron. Rev.*, 45, 449
- Lombardi, R., Gaudenzi, S. & Giovannelli, F. 1987, *Ap&SS*, 130, 275L
- Martin, J. S., Friend, M. T., Smith, R. C., & Jones, D. H. P. 1989, *MNRAS*, 240, 519
- Martinez-Pais, I. G., Giovannelli, F., Rossi, C., & Gaudenzi, S. 1994, *A&A*, 291, 455
- Mauche, C. W. 2004, *ApJ*, 610, 422
- North, R. C., Marsh, T. R., Moran, C. K. J., Kolb, U. Smith, R. C., & Stehle, R. 2000, *MNRAS*, 313, 383
- North, R. C., Marsh, T. R., Moran, C. K. J., Kolb, U. Smith, R. C., & Stehle, R. 2001, in *Astrotomography, Indirect Imaging Methods in Observational Astronomy*, Ed. by H. M. J. Boffin, D. Steeghs, & J. Cuypers, *Lect. Notes in Phys.*, 573, 33
- North, R. C., Marsh, T. R., Kolb, U., Dhillon, V. S., & Moran, C. K. J. 2002, *MNRAS*, 337, 1215

- Ritter, H., & Kolb, U. 1998, *A&AS*, 129, 83.
- Robinson, E. L. 1992, *ASP Conf Ser.*, 29, 3
- Robinson, E. L., Marsh, T. R., & Smak, J. I. 1993, in *Accretion Disks in Compact Stellar Systems*, ed. by C. Wheeler (Singapore: World Scientific Pub.), p.75.
- Robinson, E. L., Zhang, E.-H., & Stover, R. J. 1986, *ApJ*, 305, 732
- Saito, R. K., & Baptista, R. 2006, *AJ*, 131, 2185
- Shahbaz, T. 1998, *MNRAS*, 298, 153
- Smith, R. C., & Dhillon, V. S. 1998, *MNRAS*, 301, 767
- Snedden, C.A. 1973, Ph.D. Thesis, University of Texas at Austin
- Stover, R. J., Robinson, E. L., Nather, R. E., & Montemayor, T. J. 1980, *ApJ*, 240, 597
- Tokunaga, A. T. 2000, in *Allen’s Astrophysical Quantities*, ed. A.N. Cox (4th ed.;New York: Springer), 151
- Tull, R. G. 1998, *Proc. SPIE*, 3355, 387
- Voloshina, I. B. 1986, *Pis ma Astronomicheskii Zhurnal*, 12, 219
- Voloshina, I. B., & Khruzina, T. S. 2000, *Astronomy Reports*, 44, 89
- Vrielmann, S., & Baptista, R. 2002, *Astron. Nachr.*, 323, 75
- Wade, R. A. 1982, *AJ*, 87, 1558
- Walker, M. F. 1981, *ApJ*, 248, 256
- Walker, M. F., & Chincarini, G. 1968, *ApJ*, 154, 157
- Walker, M. F., & Reagan, G. H. 1971, *IBVS*, No. 544.
- Warner, B. 1995, *Cataclysmic Variable Stars* (Cambridge: Cambridge University Press)
- Warner, B. 2004, *PASP*, 116, 115
- Webb, N. A., Naylor, T., Jeffries, R. D. 2002, *ApJ*, 568, L45
- Welsh, W. F., Horne, K., & Gomer, R. 1993, *ApJ*, 410, L39

Table 1. Radial Velocities of the Secondary Star in SS Cyg

Date (HJD – 2452000)	Velocity (km s ⁻¹)	Date (HJD – 2452000)	Velocity (km s ⁻¹)
83.81229	136.2	104.77608	104.7
86.80263	31.0	105.76826	-34.6
91.80944	150.7	106.74746	-56.2
94.80816	108.8	107.78210	143.3
95.79076	-170.5	109.76639	-8.5
97.76919	-95.9	110.77827	138.2
98.80362	-155.5	111.73070	-139.7
99.78387	144.1	112.74418	122.4
100.75669	-174.9	113.72400	-110.2
101.78850	-30.6	113.76127	18.6
102.79008	121.4	113.94938	-174.8
103.78679	-168.5		

Table 2. Measurements of the Contribution of Disk Flux to the Total Flux from SS Cygni during Quiescence

Wavelength Region	Fraction of Flux from the Disk	Reference
near $H\beta$	0.50 ± 0.03	Hessman et al. (1984) quoted in Hessman (1988)
5500 Å	0.56 ± 0.03	Wade (1982)
5500 Å	0.575 ± 0.075	This work
6300 - 6500 Å	0.315 ± 0.004	North et al. (2002)
6400 - 6700 Å	0.45	Martinez-Pais et al. (1994)
7000 - 7600 Å	0.50	Webb et al. (2002)

Table 3. Previous Measurements of the Emission- and Absorption-line Radial Velocity Curves of SS Cygni

K_K (km s ⁻¹)	K_{em} (km s ⁻¹)	Reference
115	122	Joy (1956)
115	122	Walker & Chincarini (1968)
–	85 ± 4	Kiplinger (1979)
120 ± 6	118 ± 8	Cowley et al. (1980)
123 ± 2	107 ± 2	Walker (1981)
153 ± 2	90 ± 2	Stover et al. (1980)
155 ± 2	96 ± 3	Hessman et al. (1984)
152 ± 2	96 ± 5	Echevarria et al. (1989)
155 ± 3	92 ± 2	Friend et al. (1990)
162 ± 3	–	Martinez-Pais et al. (1994)
165 ± 1	–	North et al. (2002)
162.5 ± 1	–	This work

Fig. 1.— The eruption light curve of SS Cyg in 1981 and 2001 from the American Association of Variable Star Observers. The horizontal bar plotted below the light curve in 1981 marks the dates when Hessman et al. (1984) obtained their spectrograms of SS Cyg; the horizontal bar in 2001 shows when we obtained our spectrograms. Both sets of data were obtained when SS Cyg was in its quiescent state, but SS Cyg was ~ 0.6 mag fainter during quiescence in 2001 than it was in 1984.

Fig. 2.— The upper panel shows a portion of the average of all our spectrograms of SS Cyg. Before the spectrograms were averaged their orbital velocities were removed by shifting the spectrograms to SS Cyg’s systemic velocity. The lower panel shows a synthetic spectrum for a single, non-rotating star with a K4-5 V spectral type for comparison. The extra width of the absorption lines in SS Cyg is due to a combination of rotational broadening ($\sim 89 \text{ km s}^{-1}$), orbital smearing during the exposures ($< 26 \text{ km s}^{-1}$), and instrumental resolution of the spectrograph (8.2 km s^{-1}).

Fig. 3.— A contour plot of the χ^2 of the fits of the LinBrod synthetic spectra to the observed spectrum of SS Cyg as a function of K_K and q . The contours of constant χ^2 are evenly spaced multiples of the minimum χ^2 beginning at 1.0001 and separated by 0.0002. The best fit values are marked by the cross at $q = 0.685$ and $K_K = 162.5 \text{ km s}^{-1}$.

Fig. 4.— The dots and error bars are measurements of the V-band orbital light curve of SS Cyg from Voloshina & Khruzina (2000). Orbital phase $\phi = 0$ corresponds to spectroscopic conjunction with the white dwarf behind the K star. The basic double-humped variation is produced by ellipsoidal variations of the K star. The extra amplitude and the asymmetry of the hump at $\phi = 0.75$ is produced by a bright spot on the outer edge of the disk. The solid line is a fitted light curve for an orbital inclination $i = 52^\circ$ and 56% of the light at $\phi = 0.25$ coming from the accretion disk. As noted in the text and shown in Figure 5, there is a strong positive correlation between the adopted third-light fraction and the best fitting orbital inclination.

Fig. 5.— The correlation between orbital inclination and the fraction f of the V-band flux contributed by the accretion disk and white dwarf. The best fitting inclinations for any adopted fraction are shown by the solid line extending from the lower-left to the upper right in the diagram. The parallel dashed lines show one standard deviation confidence limits for the fits. The vertical solid corresponds to $f = 0.535$, derived from the dilution of the absorption lines in the spectrum of the K star; and the two vertical dashed lines correspond to $f = 0.535 \pm 0.075$, the confidence limits on f . The derived inclination lies in the interval $45^\circ \leq i \leq 56^\circ$.









